arXiv:math/0107166v3 [math.KT] 30 Apr 2002

Cyclic Cohomology of Crossed Coproduct Coalgebras

R. Akbarpour, M. Khalkhali
Department of Mathematics , University of Western Ontario
`akbarpur@uwo.ca` `masoud@uwo.ca`

February 1, 2008

Abstract

We extend our work in [1] to the case of Hopf comodule coalgebras. We introduce the cocylindrical module $C\natural\mathcal{H}$, where $\mathcal{H}$ is a Hopf algebra with bijective antipode and C is a Hopf comodule coalgebra over $\mathcal{H}$. We show that there exists an isomorphism between the cocyclic module of the crossed coproduct coalgebra $C >\!\!\blacktriangleleft \mathcal{H}$ and $\Delta(C\natural\mathcal{H})$, the cocyclic module related to the diagonal of $C\natural\mathcal{H}$. We approximate $HC^\bullet(C >\!\!\blacktriangleleft \mathcal{H})$ by a spectral sequence and we give an interpretation for $\mathsf{E}^0, \mathsf{E}^1$ and E^2 terms of this spectral sequence.

Keywords. Cyclic homology, Hopf algebras .

1 Introduction

Getzler and Jones in [5] introduced a method to compute the cyclic homology of a crossed product algebras $A \rtimes G$, where G is a group that acts on an algebra A by automorphisms. This method is based on constructing a cylindrical module, $A\natural G$, and showing that $\Delta(A\natural G) \cong \mathsf{C}_\bullet(A \rtimes G)$, where Δ is the diagonal and $\mathsf{C}_\bullet$ the cyclic module functor. Then by using the Eilenberg-Zilber theorem for cylindrical modules, they obtained a quasi-isomorphism of mixed complexes $\Delta(A\natural G) \cong Tot_\bullet(A\natural G)$, and a spectral sequence converging to $HC_\bullet(A \rtimes G)$. This spectral sequence was first obtained, by a different method, by Feigin and Tsygan [4]. We used the the same method in [1] to generalize their work to Hopf module algebras with the action of a Hopf algebra $\mathcal{H}$ on an algebra A, where the antipode of $\mathcal{H}$ is assumed to be bijective. In this paper we continue our work in [1] to obtain similar results for Hopf comodule coalgebras.

We construct a cocylindrical module $C\natural\mathcal{H}$ and we show that there exists an isomorphism between cocyclic structures of the diagonal of $C\natural\mathcal{H}$, denoted by $\Delta(C\natural\mathcal{H})$, and the crossed coproduct coalgebra $C >\!\!\blacktriangleleft \mathcal{H}$. We apply the Eilenberg-Zilber theorem for cocylindrical modules to get a quasi-isomorphism of mixed complexes, $\Delta(C\natural\mathcal{H}) \cong Tot^{\bullet}(C\natural\mathcal{H})$, and we approximate $HC^{\bullet}(C >\!\!\blacktriangleleft \mathcal{H})$ by a spectral sequence. We give an interpretation for the first three terms of this spectral sequence. In our computations we find the coinvariant cocyclic module $C^{\bullet}_{\mathcal{H}}(C)$ that is dual to $C^{\mathcal{H}}_{\bullet}(A)$ that we found in [1].

2 Preliminaries on cocylindrical modules and Hopf algebras

Let $[j], j \in \mathbb{Z}^{+}$ be the set of all integers with the standard ordering and t_j an automorphism of $[j]$ defined by $t_j(i) = i+1, i \in \mathbb{Z}$. The paracyclic category Λ_∞ is a category whose objects are $[j], j \geq 0$ and whose morphisms, $Hom_{\Lambda_\infty}([i],[j])$, are the sets of all nondecreasing maps $f : [i] \to [j]$ with $t_j^{j+1} f = f\, t_i^{i+1}$ [4].

One can identify the simplicial category Δ as a sub category of Λ_∞ so that the objects of Δ are the same as Λ_∞ and morphisms, $Hom_{\Delta}([i],[j])$, are those morphisms of $Hom_{\Lambda_\infty}([i],[j])$ that map $\{0,1,\ldots,i\}$ into $\{0,1,\ldots,j\}$. Every element $g \in Hom_{\Lambda_\infty}([i],[j])$ $,0 \leq r \leq \infty$, has a unique decomposition $g = f\, t_i^k$ where $f \in Hom_{\Delta}([i],[j])$.

If $\mathcal{A}$ is any category, a paracocyclic object in $\mathcal{A}$ is a covariant functor from Λ_∞ to $\mathcal{A}$. This definition is equivalent to giving a sequence of objects $A_0, A_1, \ldots$ together with coface operators $\partial^i : A_n \to A_{n+1}, (i = 0,1,\ldots,n+1)$, codegeneracy operators $\sigma^i : A_n \to A_{n-1}, (i = 0,1,\ldots,n-1)$, and cyclic operators $\tau_n : A_n \to A_n$ where these operators satisfy the cosimplicial and Λ_∞-identities:

$$\begin{aligned} \partial^j\partial^i &= \partial^i\partial^{j-1} \quad & i<j, \\ \sigma^j\sigma^i &= \sigma^i\sigma^{j+1} \quad & i \leq j, \end{aligned}$$

$$\sigma^j \partial^i = \begin{cases} \partial^i \sigma^{j-1} & i < j \\ \text{identity} & i = j \text{ or } i = j+1 \\ \partial^{i-1} \sigma^j & i > j+1. \end{cases}$$

$$\begin{aligned} \tau_{n+1}\partial^i &= \partial^{i-1}\tau_n, \quad 1 \leq i \leq n \ , \quad \tau_{n+1}\partial^0 = \partial^n, \\ \tau_{n-1}\sigma^i &= \sigma^{i-1}\tau_n, \quad 1 \leq i \leq n \ , \quad \tau_{n-1}\sigma^0 = \sigma^{n-1}\tau_n^2. \end{aligned} \tag{1}$$

$$\tag{2}$$

If in addition we have $\tau_n^{n+1} = id$, then we have a cocyclic object in the sense of Connes [3]. By a bi-paracocyclic object in a category $\mathcal{A}$, we mean a paracocyclic object in the category of paracocyclic objects in $\mathcal{A}$. So, giving a bi-paracocyclic object in $\mathcal{A}$ is equivalent to giving a double sequence $A(p,q)$ of objects of $\mathcal{A}$ and operators $\partial_{p,q}, \sigma_{p,q}, \tau_{p,q}$ and $\bar{\partial}_{p,q}, \bar{\sigma}_{p,q}, \bar{\tau}_{p,q}$ such that, for all $p \geq 0$,

$$B_p(q) = \{A(p,q), \sigma^i_{p,q}, \partial^i_{p,q}, \tau_{p,q}\},$$

and for all $q \geq 0$,

$$\bar{B}_q(p) = \{A(p,q), \bar{\sigma}^i_{p,q}, \bar{\partial}^i_{p,q}, \bar{\tau}_{p,q}\},$$

are paracocyclic objects in $\mathcal{A}$ and every horizontal operator commutes with every vertical operator.

We say that a bi-paracocyclic object is cocylindrical [5] if for all $p, q \geq 0$,

$$\bar{\tau}^{p+1}_{p,q} \ \tau^{q+1}_{p,q} = id_{p,q}. \tag{3}$$

If A is a bi-paracocyclic object in a category $\mathcal{A}$, the paracocyclic object related to the diagonal of A will be denoted by ΔA. So, the paracocyclic operators on $\Delta A(n) = A(n,n)$ are $\bar{\partial}^i_{n,n+1} \ \partial^i_{n,n}, \bar{\sigma}^i_{n,n-1}\sigma^i_{n,n}, \bar{\tau}_{n,n}\tau_{n,n}$. When A is a cocylindrical object since the cyclic operator of ΔA is $\bar{\tau}_{n,n}\tau_{n,n}$ and $\bar{\tau}, \tau$ commute, then, from $\bar{\tau}^{n+1}_{n,n}\tau^{n+1}_{n,n} = id_{n,n}$, we conclude that $(\bar{\tau}_{n,n}\tau_{n,n})^{n+1} = id$. So that ΔA is a cocyclic object.

Let k be a commutative unital ring. By a paracocyclic (resp. cocylindrical or cocyclic) module over k, we mean a paracocyclic (resp. cocylindrical or cocyclic) object in the category of k-modules.

An important example of a cocyclic module in this paper is the cocyclic module of a coalgebra C, denoted by $\mathsf{C}^\bullet(C)$. It is defined by $\mathsf{C}^n(C) = C^{\otimes(n+1)}, n \geq 0$, with coface, codegeneracy and cyclic oparators

$$\begin{aligned}
&\partial^i(a_0, a_1, \dots, a_n) = (a_0, \dots, a_i^{(0)}, a_i^{(1)}, \dots, a_n), \;\; 1 \leq i \leq n,\\
&\partial^{n+1}(a_0, a_1, \dots, a_n) = (a_0^{(1)}, \dots, \dots, a_n, a_0^{(0)}),\\
&\sigma^i(a_0, a_1, \dots, a_n) = \epsilon(a_i)(a_0, \dots, a_{i-1}, a_{i+1}, \dots, a_n), \;\; 1 \leq i \leq n,\\
&\tau(a_0, a_1, \dots, a_n) = (a_1, a_2, \dots, a_0).
\end{aligned}$$

A paracochain complex, by definition, is a graded k-module $\mathsf{V}^\bullet = (V^i)_{i \in \mathbb{N}}$ equipped with operators $b : V^i \to V^{i+1}$ and $B : V^i \to V^{i-1}$ such that $b^2 = B^2 = 0$, and the operator $T = 1 - (bB + Bb)$ is invertible. In the case that $T = 1$, the paracochain complex is called a mixed (cochain) complex.

Corresponding to any paracocyclic module A, we can define the paracochain complex $\mathsf{C}^\bullet(A)$ with the underlying graded module $\mathsf{C}^n(A) = A(n)$ and the operators $b = \sum_{i=0}^n (-1)^i \partial^i$ and $B = N\sigma(1 - (-1)^{n+1}\tau)$. Here, σ is the extra degeneracy satisfying $\tau\sigma^0 = \sigma\tau$, and $N = \sum_{i=0}^n (-1)^{in}\tau^i$ is the norm operator. For any bi-paracocyclic module A, $Tot(\mathsf{C}(A))$ is a paracochain complex with $Tot^n(\mathsf{C}(A)) = \sum_{p+q=n} A(p,q)$ and with the operators $Tot(b) = b + \bar{b}$ and $Tot(B) = B + T\bar{B}$, where $T = 1 - (bB + Bb)$. It is a mixed complex if A is cocylindrical [5].

If we define the normalized cochain functor N from paracocyclic modules to paracochain complexes with the underlying graded module $\mathsf{N}^n(A) = \bigcap_{i=0}^{n-1} ker(\sigma^i)$ and the operators b, B induced from $\mathsf{C}^\bullet(A)$, then we have the following well-known results(see [5] for a dual version):

1. The inclusion $(\mathsf{N}^\bullet(A), b) \to (\mathsf{C}^\bullet(A), b)$ is a quasi-isomorphism of complexes.
2. The cyclic Eilenberg-Zilber theorem holds for cocylindrical modules, i.e., for any cocylindrical module A, there is a natural quasi-isomorphism $\mathbf{f}_0 + \mathbf{u}\mathbf{f_1} : \mathsf{N}^\bullet(\Delta(A)) \to Tot^\bullet(\mathsf{N}(A))$ of mixed complexes, where $\mathbf{f}_0$ is the shuffle map.

When A is a cocyclic module, then we can construct the related mixed complex $(\mathsf{C}^\bullet(A)[\mathbf{u}], b + \mathbf{u}B)$. The cohomology of the complex $(\mathsf{C}^\bullet(A)[\mathbf{u}] \otimes_{k[\mathbf{u}]} \mathsf{W}, b + \mathbf{u}B)$ defines the cyclic cohomology of A with coefficients in W, where W is a graded $k[\mathbf{u}]$-module with finite homological dimention. We denote $\mathsf{C}_\bullet(A) \otimes_{k[\mathbf{u}]} \mathsf{W}$ by $\mathsf{C}(A) \boxtimes \mathsf{W}$. We know that if $\mathsf{W} = k[\mathbf{u}, \mathbf{u^{-1}}]$, $k[\mathbf{u}, \mathbf{u^{-1}}]/\mathbf{u}k[\mathbf{u}]$, $k[\mathbf{u}]/\mathbf{u}k[\mathbf{u}]$ we get, respectively, $HP^\bullet(A)$ the periodic cyclic cohomology, $HC^\bullet(A)$ the cyclic cohomology and $HH^\bullet(A)$ the Hochschild cohomology of A [5].

In this paper the word coalgebra means a coassociative, not necessarily cocommutative, counital coalgebra over a fixed commutative ring k. Similarly, our Hopf algebras are over k and are not assumed to be commutative or cocommutative. The undecorated tensor product $\otimes$ means tensor product over k. If $\mathcal{H}$ is a Hopf algebra, we denote its coproduct by $\Delta : \mathcal{H} \to \mathcal{H}\otimes\mathcal{H}$, its counit by $\epsilon : \mathcal{H} \to k$, its unit by $\eta : k \to \mathcal{H}$ and its antipode by $S : \mathcal{H} \to \mathcal{H}$. We will use Sweedler's notation $\Delta(h) = h^{(0)} \otimes h^{(1)}$, $\Delta^2(h) = (\Delta \otimes id) \circ \Delta(h) = (id \otimes \Delta) \circ \Delta(h) = h^{(0)} \otimes h^{(1)} \otimes h^{(2)}$, etc., where summation is understood. If $\mathcal{H}$ is a Hopf algebra, the word $\mathcal{H}$-comodule means a comodule over the underlying coalgebra of $\mathcal{H}$. We use Sweedler's notation for comodules. Thus if $\rho : M \to \mathcal{H} \otimes M$, is the structure map of a left $\mathcal{H}$-comodule M, we write $\rho(m) = m^{(\bar{1})} \otimes m^{(\bar{0})}$, where summation is understood.

A coalgebra C is called a left $\mathcal{H}$-comodule coalgebra if C is a left $\mathcal{H}$-comodule and the comultiplication map $C \to C \otimes C$ and the counit map $C \to k$ are morphisms of $\mathcal{H}$ -comodules, i.e., for all $a \in C$,

$$\sum a^{(0)(\bar{1})} a^{(1)(\bar{1})} \otimes a^{(0)(\bar{0})} \otimes a^{(1)(\bar{0})} = \sum a^{(\bar{1})} \otimes a^{(\bar{0})(0)} \otimes a^{(\bar{0})(1)},$$

$$\sum a^{(\bar{1})} \epsilon(a^{(\bar{0})}) = \epsilon(a) 1_{\mathcal{H}},$$

where ϵ is the counit of C. The reader can consult [2] for various examples of comodule coalgebras.

If $\mathcal{H}$ is a bialgebra and C a left $\mathcal{H}$-comodule coalgebra, the crossed coproduct coalgebra $C >\!\!\blacktriangleleft \mathcal{H}$ is defined as $C \otimes \mathcal{H}$ with the coalgebra structure

$$\begin{aligned} &\Delta(a \otimes g) = \sum a^{(0)} \otimes a^{(1)(\bar{1})} g^{(0)} \otimes a^{(1)(\bar{0})} \otimes g^{(1)}, \\ &\epsilon(a \otimes g) = \epsilon(a)\epsilon(g), \end{aligned}$$

for $g \in \mathcal{H}$ and $a \in C$.

For more information on cyclic homology and Hopf algebras we recommend [3, 6, 7, 8, 9].

3 The Cocylindrical Module C$\natural\mathcal{H}$

Let $\mathcal{H}$ be a Hopf algebra with bijective antipode and C a left $\mathcal{H}$-comodule coalgebra. In this section we associate a cocylindrical module $C\natural\mathcal{H}$ to this data by $C\natural\mathcal{H} = \{\mathcal{H}^{\otimes(p+1)} \otimes C^{\otimes(q+1)}\}_{p,q\geq 0}$. We define the operators $\tau_{p,q}, \partial_{p,q}, \sigma_{p,q}$ and $\bar{\tau}_{p,q}, \bar{\partial}_{p,q}, \bar{\sigma}_{p,q}$ as follows:

$$\tau_{p,q}(g_0, \dots, g_p \mid a_0, \dots, a_q) = (S^{-1}(a_0^{(\bar{1})}) \cdot (g_0, \dots, g_p) \mid a_1, \dots, a_q, a_0^{(\bar{0})}),$$

$$\begin{aligned}
&\sigma_{p,q}^{i}(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= (g_0, \dots, g_p \mid a_0, \dots, a_i, a_{i+2}, \dots, a_q)\epsilon(a_{i+1}) \quad 0 \leq i < q,\\
&\partial_{p,q}^{i}(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= (g_0, \dots, g_p \mid a_0, \dots, a_i^{(0)}, a_i^{(1)}, a_{i+1}, \dots, a_q) \quad 0 \leq i \leq q,\\
&\partial_{p,q}^{q+1}(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= (S^{-1}(a_0^{(0)(\bar{1})}) \cdot (g_1, \dots, g_p) \mid a_0^{(1)}, a_0, \dots, a_q, a_0^{(0)(\bar{0})}),
\end{aligned} \tag{4}$$

where $\quad h \cdot (g_0, \dots, g_p) := (h^{(0)}g_0, \dots, h^{(p)}g_p)$, and we see that

$$\begin{aligned}
&S^{-1}(a_0^{(\bar{1})}) \cdot (g_0, \dots, g_p) = (S^{-1}(a_0^{(\bar{1})(p)})g_0, \dots, S^{-1}(a_0^{(\bar{1})(0)})g_p)\\
= \quad &(S^{-1}(a_0^{(\bar{1})})g_0, \dots, S^{-1}(a_0^{(\overline{p+1})})g_p).
\end{aligned}$$

$$\begin{aligned}
&\bar{\tau}_{p,q}(g_0, \dots, g_p \mid a_0, \dots, a_q) = (g_1, \dots, g_{p-1}, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_0 \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})}),\\
&\bar{\sigma}_{p,q}^{i}(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= (g_0, \dots, g_i, g_{i+2}, \dots, , g_p \mid a_0, \dots, a_q)\epsilon(g_{i+1}) \;\; 0 \leq i < p,\\
&\bar{\partial}_{p,q}^{i}(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= (g_0, \dots, g_i^{(0)}, g_i^{(1)}, \dots, g_p \mid a_0, \dots, a_q) \quad 0 \leq i \leq p,\\
&\bar{\partial}_{p,q}^{p+1}(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= (g_0^{(1)}, g_0, \dots, g_p, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_0^{(0)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})}).
\end{aligned} \tag{5}$$

Theorem 3.1. *$C\natural\mathcal{H}$ with the operators defined in (4),(5) is a cocylindrical module.*

Proof. We verify only some of the important identities. The other identities are similar to check.

- $\tau_{p,q+1}\partial^0_{p,q} = \partial^{q+1}_{p,q}$

  $\tau_{p,q+1}\partial^0_{p,q}(g_0, \dots, g_p \mid a_0, \dots, a_q)$

$$\begin{aligned} &= \tau_{p,q+1}(g_0, \dots, g_p \mid a_0^{(0)}, a_0^{(1)}, \dots, a_q) \\ &= \partial_{p,q}(S^{-1}(a_0^{(0)(\bar{1})}) \cdot (g_0, \dots, g_p) \mid a_0^{(1)}, \dots, a_q, a_0^{(0)(\bar{0})}) \\ &= \partial^{q+1}_{p,q}(g_0, \dots, g_p \mid a_0, \dots, a_q). \end{aligned}$$

- $\tau_{p,q-1}\sigma^0_{p,q} = \sigma^{q+1}_{p,q}(\tau^q_{p,q})^2$

$$\tau_{p,q-1}\sigma^0_{p,q}(g_0, \dots, g_p \mid a_0, \dots, a_q) = (S^{-1}(a_0^{(\bar{1})}) \cdot (g_0, \dots, g_p) \mid a_1, \dots, a_q, a_0^{(\bar{0})})\epsilon(a_1),$$

$$\sigma^{q+1}_{p,q}(\tau_{p,q})^2(g_0, \dots, g_p \mid a_0, \dots, a_q)$$

$$\begin{aligned} &= \sigma^{q+1}_{p,q}((S^{-1}(a_1^{(\bar{1}))})(S^{-1}(a_0^{(\bar{1}))})) \cdot (g_0, \dots, g_p) \mid a_2, \dots, a_q, a_0^{(\bar{0})}, a_1^{(\bar{0})}) \\ &= (S^{-1}(a_0^{(\bar{1})}) \cdot (g_0, \dots, g_p) \mid a_1, \dots, a_q, a_0^{(\bar{0})})\epsilon(a_1). \end{aligned}$$

- $\bar{\tau}_{p-1,q}\bar{\sigma}^0_{p,q} = \bar{\sigma}^{p-1}_{p,q}(\bar{\tau}_{p,q})^2$

$$\bar{\tau}_{p-1,q}\bar{\sigma}^0_{p,q}(g_0, \dots, g_p \mid a_0, \dots, a_q) = (g_2, \dots, g_p, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_0 \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\epsilon(g_1),$$

$$\bar{\sigma}^{p-1}_{p,q}(\bar{\tau}_{p,q})^2(g_0, \dots, g_p \mid a_0, \dots, a_q)$$

$$\begin{aligned} &= \bar{\sigma}^{p-1}_{p,q}(g_2, \dots, g_p, (a_0^{(\bar{2})} \dots a_q^{(\bar{2})})g_0, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_1 \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})}) \\ &= (g_2, \dots, g_p, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_0 \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\epsilon(g_1). \end{aligned}$$

- $\bar{\tau}_{p+1,q}\bar{\partial}^0_{p,q} = \bar{\partial}^{q+1}_{p,q}$

  $\bar{\tau}_{p+1,q}\bar{\partial}^0_{p,q}(g_0, \dots, g_p \mid a_0, \dots, a_q)$

$$\begin{aligned} &= (g_0^{(1)}, g_1, \dots, g_{p-1}, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_0^{(0)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})}) \\ &= \bar{\partial}^{p,q}_{q+1}(g_0, \dots, g_p \mid a_0, \dots, a_q). \end{aligned}$$

- $\tau_{p,q}\bar{\tau}_{p,q} = \bar{\tau}_{p,q}\tau_{p,q}$

$$\begin{aligned}
&(\tau_{p,q}\bar{\tau}_{p,q})(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= \tau_{p,q}(g_1, \dots, g_p, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_0 \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\\
&= (S^{-1}(a_0^{(\bar{1})}) \cdot (g_1, \dots g_p, (a_0^{(\bar{2})} \dots a_q^{(\bar{1})})g_0) \mid a_1^{(\bar{0})}, \dots a_q^{(\bar{0})}, a_0^{(\bar{0})})\\
&= (S^{-1}(a_0^{(\bar{1})(0)}) \cdot (g_1, \dots g_p), S^{-1}(a_0^{(\bar{1})(1)})(a_0^{(\bar{1})(2)} a_1^{(\bar{1})} \dots a_q^{(\bar{1})})g_0) \mid a_1^{(\bar{0})}, \dots a_q^{(\bar{0})}, a_0^{(\bar{0})})\\
&= (S^{-1}(a_0^{(\bar{1})}) \cdot (g_1, \dots g_p), (a_1^{(\bar{1})} \dots a_q^{(\bar{1})})g_0 \mid a_1^{(\bar{0})}, \dots a_q^{(\bar{0})}, a_0^{(\bar{0})}).
\end{aligned}$$

$$\begin{aligned}
&(\bar{\tau}_{p,q}\tau_{p,q})(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= \bar{\tau}_{p,q}(S^{-1}(a_0^{(\bar{1})}) \cdot (g_0, \dots, g_p) \mid a_1, \dots, a_q, a_0^{(\bar{0})})\\
&= \bar{\tau}_{p,q}(S^{-1}(a_0^{(\bar{1})(0)})g_0, S^{-1}(a_0^{(\bar{1})(1)}) \cdot (g_1, \dots, g_p) \mid a_1, \dots, a_q, a_0^{(\bar{0})})\\
&= (S^{-1}(a_0^{(\bar{2})(1)}) \cdot (g_1, \dots, g_p), (a_1^{(\bar{1})} \dots a_p^{(\bar{1})} a_0^{(\bar{1})})S^{-1}(a_0^{(\bar{2})(1)})g_0) \mid a_1, \dots, a_q, a_0^{(\bar{0})})\\
&= (S^{-1}(a_0^{(\bar{1})(1)}) \cdot (g_1, \dots, g_p), (a_1^{(\bar{1})} \dots a_p^{(\bar{1})} a_0^{(\bar{1})(0)})S^{-1}(a_0^{(\bar{1})(1)})g_0) \mid a_1, \dots, a_q, a_0^{(\bar{0})})\\
&= (S^{-1}(a_0^{(\bar{1})}) \cdot (g_1, \dots, g_p), (a_1^{(\bar{1})} \dots a_p^{(\bar{1})})g_0) \mid a_1, \dots, a_q, a_0^{(\bar{0})}).
\end{aligned}$$

- $\bar{\tau}_{p,q}^{p+1}\ \tau_{p,q}^{q+1} = \tau_{p,q}^{q+1}\ \bar{\tau}_{p,q}^{p+1} = id_{p,q}$

$$\begin{aligned}
&\tau_{p,q}^{q+1}\bar{\tau}_{p,q}^{p+1}(g_0, \dots, g_p \mid a_0, \dots, a_q)\\
&= \tau_{p,q}^{q+1}((a_0^{(\overline{p+1})} \dots a_q^{(\overline{p+1})})g_0, \dots, (a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g_p \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\\
&= S^{-1}(a_0^{(\bar{1})} \dots a_q^{(\bar{1})}) \cdot ((a_0^{(\overline{p+2})} \dots a_q^{(\overline{p+2})})g_0, \dots, (a_0^{(\bar{2})} \dots a_q^{(\bar{2})})g_p) \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\\
&= (g_0, \dots, g_p \mid a_0, \dots, a_q).
\end{aligned}$$

□

Corollary 3.1. *The diagonal $\Delta(C\natural\mathcal{H})$ is a cocyclic module.*

4 Relation of $\Delta(\mathbf{C}\natural\mathcal{H})$ with the Cocyclic Module of the Crossed Coproduct Coalgebra $\mathbf{C} >\!\!\blacktriangleleft \mathcal{H}$

We define a map $\phi : \mathsf{C}^\bullet(C >\!\!\blacktriangleleft \mathcal{H}) \to \Delta(C\natural\mathcal{H})$, by

$$\phi(a_0 \otimes g_0, \dots, a_n \otimes g_n) =$$

$$(S^{-1}(a_1^{(\bar{1})}\dots a_p^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_p^{(\bar{2})})g_1, \dots, S^{-1}(a_p^{(\overline{p})})g_{p-1}, g_p \mid a_0, a_1^{(\bar{0})}, \dots, a_q^{(\bar{0})}),$$

where $\mathsf{C}^\bullet(C >\!\!\blacktriangleleft \mathcal{H})$ denotes the cocyclic module of the crossed coproduct coalgebra $C >\!\!\blacktriangleleft \mathcal{H}$.

Theorem 4.1. *$\phi : \mathsf{C}^\bullet(C >\!\!\blacktriangleleft \mathcal{H}) \to \Delta(C\natural\mathcal{H})$ defines a cocyclic map.*

Proof. We show that ϕ commutes with the cosimplicial and cyclic operators:

- $\tau_{n,n}\bar{\tau}_{n,n}\phi = \phi\tau_n^{C>\!\!\blacktriangleleft\mathcal{H}}$

$$\begin{aligned}
& \tau_{n,n}\bar{\tau}_{n,n}\phi(a_0 \otimes g_0, \dots, a_n \otimes g_n) \\
= \;& \tau_{n,n}\bar{\tau}_{n,n}(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1, \dots, S^{-1}(a_n^{(\overline{n})})g_{n-1}, g_n \mid a_0, a_1^{(\bar{0})}, \dots, a_n^{(\bar{0})}) \\
= \;& \tau_{n,n}(S^{-1}(a_2^{(\bar{3})}\dots a_n^{(\bar{3})})g_1, \dots, S^{-1}(a_n^{(\overline{n+2})})g_n, (a_0^{(\bar{1})}\dots a_n^{(\bar{1})})S^{-1}(a_1^{(\bar{2})}\dots a_n^{(\bar{2})})g_0 \\
& \qquad\qquad \mid a_0^{(\bar{0})}, a_1, \dots, a_n^{(\bar{0})}) \\
= \;& \tau_{n,n}(S^{-1}(a_2^{(\bar{1})}\dots a_n^{(\bar{1})})g_1, \dots, g_n, a_0^{(\bar{1})}g_0 \mid a_0^{(\bar{0})}, \dots, a_n^{(\bar{0})}) \\
= \;& (S^{-1}(a_0^{(\bar{1})}) \cdot (S^{-1}(a_2^{(\bar{1})}\dots a_n^{(\bar{1})})g_1, \dots, g_n, a_0^{(\bar{2})}g_0) \mid a_1, a_2^{(\bar{0})}, \dots, a_n^{(\bar{0})}, a_0^{(\bar{0})}) \\
= \;& (S^{-1}(a_2^{(\bar{1})}\dots a_n^{(\bar{1})}a_0^{(\bar{1})})g_1, \dots, S^{-1}(a_0^{(\overline{n})})g_n, S^{-1}(a_0^{(\overline{n+1})})a_0^{(\overline{n+2})}g_0) \mid a_1, a_2^{(\bar{0})}, \dots, a_n^{(\bar{0})}, a_0^{(\bar{0})}) \\
= \;& (S^{-1}(a_2^{(\bar{1})}\dots a_n^{(\bar{1})}a_0^{(\bar{1})})g_1, \dots, S^{-1}(a_0^{(\overline{n})})g_n, g_0) \mid a_1, a_2^{(\bar{0})}, \dots, a_n^{(\bar{0})}, a_0^{(\bar{0})}) \\
= \;& \phi(a_1 \otimes g_1, \dots, a_n \otimes g_n, a_0 \otimes g_0) = \phi\tau_n^{C>\!\!\blacktriangleleft\mathcal{H}}(a_0 \otimes g_0, \dots, a_n \otimes g_n).
\end{aligned}$$

- $\sigma_{n,n}^i\bar{\sigma}_{n,n}^i\phi = \phi\sigma_i^{C>\!\!\blacktriangleleft\mathcal{H}} \quad 0 \le i < n$

$$\begin{aligned}
& \sigma_{n,n}^i\bar{\sigma}_{n,n}^i\phi(a_0 \otimes g_0, \dots, a_n \otimes g_n) \\
= \;& \sigma_{n,n}^i\bar{\sigma}_{n,n}^i(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1, \dots, S^{-1}(a_n^{(\overline{n})})g_{n-1}, g_n \mid a_0, a_1^{(\bar{0})}, \dots, a_n^{(\bar{0})}) \\
= \;& \sigma_{n,n}^i(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1, \dots, S^{-1}(a_n^{(\overline{n})})g_{n-1}, g_n \\
& \qquad\qquad \mid a_0, a_1^{(\bar{0})}, \dots, a_n^{(\bar{0})})\epsilon(S^{-1}(a_{i+2}^{(\overline{i+2})}\dots a_n^{(\overline{i+2})})g_{i+1}) \\
= \;& (S^{-1}(a_1^{(\bar{1})}\dots a_{i-1}^{(\bar{1})}a_{i+1}^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_{i-1}^{(\bar{2})}a_{i+1}^{(\bar{2})}\dots a_n^{(\bar{2})})g_1, \dots, S^{-1}(a_n^{(\overline{n})})g_{n-1}, g_n \\
& \qquad\qquad \mid a_0, a_1^{(\bar{0})}, \dots, a_i^{(\bar{0})}, a_{i+2}^{(\bar{0})}, \dots, a_n^{(\bar{0})})\epsilon(g_{i+1})\epsilon(a_{i+1}) \\
= \;& (S^{-1}(a_1^{(\bar{1})}\dots a_{i-1}^{(\bar{1})}a_{i+1}^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_{i-1}^{(\bar{2})}a_{i+1}^{(\bar{2})}\dots a_n^{(\bar{2})})g_1, \dots, S^{-1}(a_n^{(\overline{n})})g_{n-1}, g_n \\
& \qquad\qquad \mid a_0, a_1^{(\bar{0})}, \dots, a_i^{(\bar{0})}, a_{i+2}^{(\bar{0})}, \dots, a_n^{(\bar{0})})\epsilon(g_{i+1} \otimes a_{i+1}) \\
= \;& \phi\sigma_i^{C>\!\!\blacktriangleleft\mathcal{H}}(a_0 \otimes g_0, \dots, a_n \otimes g_n).
\end{aligned}$$

- $\partial^{n+1}_{n,n}\bar{\partial}^{n+1}_{n,n}\phi = \phi\partial^{C>\!\!\blacktriangleleft\mathcal{H}}_{n+1}$

$$\begin{aligned}
&\partial^{n+1}_{n,n}\bar{\partial}^{n+1}_{n,n}\phi(a_0\otimes g_0,\dots,a_n\otimes g_n)\\
=\;&\partial^{n+1}_{n,n}\bar{\partial}^{n+1}_{n,n}(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,S^{-1}(a_n^{(\overline{n})})g_{n-1},g_n\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\mid a_0,a_1^{(\bar{0})},\dots,a_n^{(\bar{0})})\\
\\
=\;&\partial^{n+1}_{n,n}(S^{-1}(a_1^{(\bar{2})(0)}\dots a_n^{(\bar{2})(0)})g_0^{(1)}, S^{-1}(a_2^{(\bar{3})}\dots a_n^{(\bar{3})})g_1,\dots,g_n,\\
&\qquad\qquad\qquad(a_0^{(\bar{1})}\dots a_n^{(\bar{1})})S^{-1}(a_1^{(\bar{2})(1)}\dots a_n^{(\bar{2})(1)})g_0^{(0)}\mid a_0^{(\bar{0})},\dots,a_n^{(\bar{0})})\\
=\;&\partial^{n+1}_{n,n}(S^{-1}(a_0^{(\bar{3})}\dots a_n^{(\bar{3})})g_0^{(1)}, S^{-1}(a_1^{(\bar{4})}\dots a_n^{(\bar{4})})g_1,\dots,g_n,\\
&\qquad\qquad\qquad(a_0^{(\bar{1})}\dots a_n^{(\bar{1})})S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_0^{(0)}\mid a_0^{(\bar{0})},\dots,a_n^{(\bar{0})})\\
=\;&\partial^{n+1}_{n,n}(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0^{(1)}, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,g_n,a_0^{(\bar{1})}g_0^{(0)}\mid a_0^{(\bar{0})},\dots,a_n^{(\bar{0})})\\
\\
=\;&(S^{-1}(a_0^{(0)(\bar{1})})\cdot(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0^{(1)}, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,g_n,a_0^{(0)(\bar{2})}a_0^{(1)(\bar{1})}g_0^{(0)})\\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\mid a_0^{(1)(\bar{0})},\dots,a_n^{(\bar{0})},a_0^{(0)(\bar{0})})\\
=\;&(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})}a_0^{(0)(\bar{1})})g_0^{(0)}, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})}a_0^{(0)(\bar{2})})g_1,\dots,S^{-1}(a_0^{(0)(\overline{n+1})})g_n,\\
&\qquad\qquad S^{-1}(a_0^{(0)(\overline{n+2})})a_0^{(0)(\overline{n+3})}a_0^{(1)(\bar{1})}g_0^{(0)})\mid a_0^{(1)(\bar{0})},\dots,a_n^{(\bar{0})},a_0^{(0)(\bar{0})})\\
=\;&(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})}a_0^{(0)(\bar{1})})g_0^{(1)}, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})}a_0^{(0)(\bar{2})})g_1,\dots,S^{-1}(a_0^{(0)(\overline{n+1})})g_n,\\
&\qquad\qquad\qquad\qquad\qquad\qquad a_0^{(1)(\bar{1})}g_0^{(0)})\mid a_0^{(1)(\bar{0})},\dots,a_n^{(\bar{0})},a_0^{(0)(\bar{0})}).
\end{aligned}$$

$$\begin{aligned}
&\phi\partial^{C>\!\!\blacktriangleleft\mathcal{H}}_{n+1}(a_0\otimes g_0,\dots,a_n\otimes g_n)\\
&\quad=\;\phi(a_0^{(1)(\bar{0})}\otimes g_0^{(1)},a_1\otimes g_1,\dots,a_n\otimes g_n,a_0^{(0)}\otimes a_0^{(1)(\bar{1})}g_0^{(0)})\\
&\quad=\;(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})}a_0^{(0)(\bar{1})})g_0^{(1)}, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})}a_0^{(0)(\bar{2})})g_1,\dots,\\
&\qquad\qquad\qquad S^{-1}(a_0^{(0)(\overline{n+1})})g_n,a_0^{(1)(\bar{1})}g_0^{(0)})\mid a_0^{(1)(\bar{0})},\dots,a_n^{(\bar{0})},a_0^{(0)(\bar{0})})
\end{aligned}$$

- $\partial^i_{n,n}\bar{\partial}^i_{n,n}\phi = \phi\partial_i^{C\mathbin{>\!\!\blacktriangleleft}\mathcal{H}} \quad 0 \le i \le n$

$$\begin{aligned}
&\partial^i_{n,n}\bar{\partial}^i_{n,n}\phi(a_0\otimes g_0,\dots,a_n\otimes g_n)\\
=\;&\partial^i_{n,n}\bar{\partial}^i_{n,n}(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,S^{-1}(a_n^{(\overline{n})})g_{n-1},g_n \mid a_0,a_1^{(\bar{0})},\dots,a_n^{(\bar{0})})\\
=\;&\partial^i_{n,n}(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,S^{-1}(a_{i+1}^{(\overline{i+1})(1)}\dots a_{i+1}^{(\overline{i+1})(1)})g_i^{(0)},\\
&\qquad S^{-1}(a_{i+1}^{(\overline{i+1})(0)}\dots a_{i+1}^{(\overline{i+1})(0)})g_i^{(1)},\dots,S^{-1}(a_n^{(\overline{n})})g_{n-1},g_n \mid a_0,a_1^{(\bar{0})},\dots,a_n^{(\bar{0})})\\
=\;&\partial^i_{n,n}(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,S^{-1}(a_{i+1}^{(\overline{i+1})}\dots a_n^{(\overline{i+1})})g_i^{(0)},\\
&\qquad S^{-1}(a_{i+1}^{(\overline{i+2})}\dots a_n^{(\overline{i+2})})g_i^{(1)},\dots,S^{-1}(a_n^{(\overline{n+1})})g_{n-1},g_n \mid a_0,a_1^{(\bar{0})},\dots,a_n^{(\bar{0})})\\
=\;&(S^{-1}(a_1^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,S^{-1}(a_{i+1}^{(\overline{i+1})}\dots a_n^{(\overline{i+1})})g_i^{(0)},\\
&\; S^{-1}(a_{i+1}^{(\overline{i+2})}\dots a_n^{(\overline{i+2})})g_i^{(1)},\dots,S^{-1}(a_n^{(\overline{n+1})})g_{n-1},g_n \mid a_0,a_1^{(\bar{0})},\dots,a_i^{(\bar{0})(0)},a_i^{(\bar{0})(1)},\dots,a_n^{(\bar{0})}).
\end{aligned}$$

$$\begin{aligned}
&\phi\partial_i^{C\mathbin{>\!\!\blacktriangleleft}\mathcal{H}}(a_0\otimes g_0,\dots,a_n\otimes g_n)\\
=\;&\phi(a_0\otimes g_0,\dots,a_i^{(0)}\otimes a_i^{(1)(\bar{1})}g_i^{(0)},a_i^{(1)(\bar{0})}\otimes g_i^{(1)},\dots,a_n\otimes g_n)\\
=\;&(S^{-1}(a_1^{(\bar{1})}\dots a_i^{(0)(\bar{1})}a_i^{(1)(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_i^{(0)(\bar{2})}a_i^{(1)(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,\\
&\; S^{-1}(a_i^{(0)(\bar{i})}a_i^{(1)(\bar{i})}\dots a_n^{(\bar{i})})g_{i-1}, S^{-1}(a_i^{(1)(\overline{i+1})}\dots a_n^{(\overline{i+1})})a_i^{(1)(\overline{i+2})}g_i^{(0)},\\
&\quad S^{-1}(a_{i+1}^{(\overline{i+2})}\dots a_n^{(\overline{i+2})})g_i^{(1)},\dots,S^{-1}(a_n^{(\overline{n+1})})g_{n-1},g_n \mid a_0,a_1^{(\bar{0})},\dots,a_i^{(0)(\bar{0})},a_i^{(1)(\bar{0})},\dots a_n^{(\bar{0})})\\
=\;&(S^{-1}(a_1^{(\bar{1})}\dots a_i^{(\bar{1})}\dots a_n^{(\bar{1})})g_0, S^{-1}(a_2^{(\bar{2})}\dots a_i^{(\bar{2})}\dots a_n^{(\bar{2})})g_1,\dots,S^{-1}(a_i^{(\bar{i})}\dots a_n^{(\bar{i})})g_{i-1}\\
&\qquad ,S^{-1}(a_{i+1}^{(\overline{i+1})}\dots a_n^{(\overline{i+1})})g_i^{(0)},S^{-1}(a_{i+1}^{(\overline{i+2})}\dots a_n^{(\overline{i+2})})g_i^{(1)},\dots,S^{-1}(a_n^{(\overline{n+1})})g_{n-1},g_n\\
&\qquad\qquad\qquad \mid a_0,a_1^{(\bar{0})},\dots,a_i^{(\bar{0})(0)},a_i^{(\bar{0})(1)},\dots a_n^{(\bar{0})}).
\end{aligned}$$

□

Theorem 4.2. *We have an isomorphism of cocyclic modules*

$$\Delta(C\natural\mathcal{H}) \cong \mathsf{C}^\bullet(C\mathbin{>\!\!\blacktriangleleft}\mathcal{H}).$$

Proof. We define a map $\psi : \Delta(C\natural\mathcal{H}) \to \mathsf{C}^\bullet(C\mathbin{>\!\!\blacktriangleleft}\mathcal{H})$, by

$$\begin{aligned}
&\psi(g_0,\dots,g_n \mid a_0,\dots,a_n) =\\
&(a_0\otimes(a_1^{(\bar{1})}\dots a_p^{(\bar{p})})g_0, a_1^{(\bar{0})}\otimes(a_2^{(\bar{2})}\dots a_p^{(\overline{p-1})})g_1,\dots,a_{p-1}^{(\bar{0})}\otimes(a_p^{(\bar{0})})g_{p-1},a_p^{(\bar{0})}\otimes g_p).
\end{aligned}$$

One can check that ψ is a cocyclic map and $\phi \circ \psi = \psi \circ \phi = id$. For example $\phi \circ \psi(g_0, \dots, g_n \mid a_0, \dots, a_n) =$

$$
\begin{aligned}
&(S^{-1}(a_1^{(\bar{1})} \dots a_p^{(\bar{1})})(a_1^{(\bar{2})} a_2^{(\bar{4})} \dots a_p^{(\overline{2p})})g_0, S^{-1}(a_2^{(\bar{2})} \dots a_p^{(\bar{2})})(a_2^{(\bar{3})} a_3^{(\bar{5})} \dots a_p^{(\overline{2p-1})})g_1, \\
&\qquad\qquad \dots, S^{-1}(a_p^{(\overline{p})})a_p^{(\overline{p+1})} g_{p-1}, g_p \mid a_0, a_1^{(\bar{0})}, \dots, a_p^{(\bar{0})}) \\
&= (g_0, \dots, g_n \mid a_0, \dots, a_n).
\end{aligned}
$$

□

Now, we can use the Eilenberg-Zilber theorem for cocylindrical modules to conclude

$$H^\bullet(Tot(C\natural\mathcal{H}); \mathsf{W}) \cong HC^\bullet(\Delta(C\natural\mathcal{H}); \mathsf{W}) \cong HC^\bullet(C >\!\!\blacktriangleleft \mathcal{H}; \mathsf{W}).$$

5 The Cocyclic Module $\mathsf{C}^\bullet_{\mathcal{H}}(C)$ of Coinvariants and Cyclic Cohomology of $C >\!\!\blacktriangleleft \mathcal{H}$

Let M be a left $\mathcal{H}$-comodule with the coaction Δ_M. We define the space of p-cochains on $\mathcal{H}$ with values in M by $\mathsf{C}^p(\mathcal{H}, M) = \mathcal{H}^{\otimes p} \otimes M$, and we define the coboundary $\boldsymbol{\delta} : \mathsf{C}^p(\mathcal{H}, M) \to \mathsf{C}^{p+1}(\mathcal{H}, M)$ by

$$
\begin{aligned}
&\boldsymbol{\delta}(g_1, g_2, \dots, g_p, m) = (1, g_1, \dots, g_p, m) + \\
&\sum_{i=1}^{p} (-1)^i (g_1, \dots, g_i^{(0)}, g_i^{(1)}, \dots, g_p, m) + (-1)^{p+1}(g_1, \dots, g_p, \Delta_M(m)).
\end{aligned}
\tag{6}
$$

We denote the p-th cohomology of the complex $(\mathsf{C}^\bullet(\mathcal{H}, M), \boldsymbol{\delta})$ by $H^p(\mathcal{H}, M)$.

Now we define a left $\mathcal{H}$-coaction on the first row of $C\natural\mathcal{H}$, $C^\natural_{\mathcal{H}} = \{\mathcal{H} \otimes C^{\otimes(n+1)}\}_{n \geq 0}$, by

$$\boldsymbol{\Delta}(g \mid a_0, \dots, a_n) = (S^{-1}(g^{(2)})(a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g^{(0)} \mid g^{(1)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})}). \tag{7}$$

We define $\mathsf{C}^q(C^\natural_{\mathcal{H}}) = \mathcal{H}^{\otimes q} \otimes C^\natural_{\mathcal{H}}$ and a coaction on it by $\boldsymbol{\Delta}(g_1, \dots, g_p \mid m) = (g_1, \dots, g_p \mid \boldsymbol{\Delta}(m))$ where $m \in C^\natural_{\mathcal{H}}$. So we can construct $H^p(\mathcal{H}, \mathsf{C}^q(C^\natural_{\mathcal{H}}))$. We define $\mathsf{C}^\bullet_{\mathcal{H}}(C)$ as the coinvariant subspace of $C^\natural_{\mathcal{H}}$ under the above coaction, so that $\mathsf{C}^n_{\mathcal{H}}(C)$ is the space of all $(g \mid a_0, \ldots, a_n)$ such that

$$\boldsymbol{\Delta}(g \mid a_0, \ldots, a_n) = (1 \mid g \mid a_0, \ldots, a_n).$$

Now we replace the complex $(C\natural\mathcal{H}, (\partial, \sigma, \tau), (\bar{\partial}, \bar{\sigma}, \bar{\tau}))$ with the complex $(\mathsf{C}^p(\mathcal{H}, \mathsf{C}^q(C^\natural_{\mathcal{H}}), (\mathfrak{d}, \mathfrak{s}, \mathfrak{t}), (\bar{\mathfrak{d}}, \bar{\mathfrak{s}}, \bar{\mathfrak{t}}))$, under the transformation defined by the maps $\boldsymbol{\beta} : (C\natural\mathcal{H})_{p,q} \to \mathsf{C}^p(\mathcal{H}, \mathsf{C}^q(C^\natural_{\mathcal{H}}))$, and $\boldsymbol{\gamma} : \mathsf{C}^p(\mathcal{H}, \mathsf{C}^q(C^\natural_{\mathcal{H}})) \to (C\natural\mathcal{H})_{p,q}$, defined by

$$\begin{aligned}
&\boldsymbol{\beta}(g_0, \ldots, g_p \mid a_0, \ldots, a_q) = (S^{-1}(g^{(1)}) \cdot (g_1, \ldots, g_p) \mid g_0^{(0)} \mid a_0, \ldots, a_q),\\
&\boldsymbol{\gamma}(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = (g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p) \mid a_0, \ldots, a_q).
\end{aligned}$$

One can check that $\boldsymbol{\beta} \circ \boldsymbol{\gamma} = \boldsymbol{\gamma} \circ \boldsymbol{\beta} = id$. We find the operators $(\mathfrak{d}, \mathfrak{s}, \mathfrak{t})$ and $(\bar{\mathfrak{d}}, \bar{\mathfrak{s}}, \bar{\mathfrak{t}})$ under this transformation.

- First we compute $\bar{\mathfrak{b}} = \boldsymbol{\beta}\bar{b}\boldsymbol{\gamma}$. Since

$$\begin{aligned}
\bar{b}(g_0, \ldots, g_p \mid a_0, \ldots, a_q) \;&=\; \sum_{0 \le i \le p} (-1)^i (g_0, \ldots, g_i^{(0)}, g_i^{(1)}, \ldots, g_p \mid a_0, \ldots, a_q)\\
&+\; (-1)^{(p+1)} (g_0^{(1)}, \ldots, g_p, (a_q^{(\bar{1})} \ldots a_0^{(\bar{1})}) g_0^{(0)}) \mid a_0^{(\bar{0})}, \ldots, a_q^{(\bar{0})}),
\end{aligned}$$

we see that,

$$\begin{aligned}
&\bar{\mathfrak{b}}(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = \boldsymbol{\beta}\bar{b}(g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p) \mid a_0, \ldots, a_q)\\
=\;& \boldsymbol{\beta}\{(g^{(0)}, g^{(1)}, g^{(2)} \cdot (g_1, \ldots, g_p) \mid a_0, \ldots, a_q)\\
+\;& \sum_{1 \le i \le p} (-1)^i (g^{(0)}, g^{(1)} g_1, \ldots, g^{(i)} g_i^{(0)}, g^{(i+1)} g_i^{(1)}, \ldots, g^{(p+1)} g_p \mid a_0, \ldots, a_q)\\
+\;& (-1)^{(p+1)} (g^{(1)}, g^{(2)} \cdot (g_1, \ldots, g_p), (a_0^{(\bar{1})} \ldots a_q^{(\bar{1})}) g^{(0)} \mid a_0^{(\bar{0})}, \ldots, a_q^{(\bar{0})})\\
=\;& (S^{-1}(g^{(1)}) \cdot (g^{(2)}, g^{(3)} \cdot (g_1, \ldots, g_p)) \mid g^{(0)} \mid a_0, \ldots, a_q)\\
+\;& \sum_{1 \le i \le p} (-1)^i (S^{-1}(g^{(1)}) \cdot (g^{(2)} g_1, \ldots, g^{(i+1)} g_i^{(0)}, g^{(i+2)} g_i^{(1)}, \ldots, g^{(p+2)} g_p) \mid g^{(0)} \mid a_0, \ldots, a_q)\\
+\;& (-1)^{(p+1)} (S^{-1}(g^{(2)}) \cdot (g^{(3)} \cdot (g_1, \ldots, g_p), (a_0^{(\bar{1})} \ldots a_q^{(\bar{1})}) g^{(0)}) \mid g^{(1)} \mid a_0^{(\bar{0})}, \ldots, a_q^{(\bar{0})})\\
=\;& (S^{-1}(g^{(2)}) g^{(3)}, (S^{-1}(g^{(1)}) g^{(4)}) \cdot (g_1, \ldots, g_p) \mid g^{(0)} \mid a_0, \ldots, a_q)
\end{aligned}$$

$$\begin{aligned}
&+ \sum_{1\leq i\leq p} (-1)^i (S^{-1}(g^{(1)})\cdot(g^{(2)}\cdot(g_1,\ldots,g_i^{(0)},g_i^{(1)},\ldots,g_p)) \mid g^{(0)} \mid a_0,\ldots,a_q)\\
&+ (-1)^{(p+1)}((S^{-1}(g^{(3)})(g^{(4)})\cdot(g_1,\ldots,g_p), S^{-1}(g^{(2)})(a_0^{(\bar{1})}\ldots a_q^{(\bar{1})})g^{(0)}) \mid g^{(1)} \mid a_0^{(\bar{0})},\ldots,a_q^{(\bar{0})})\\
&= (1,g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q)\\
&+ \sum_{1\leq i\leq p} (-1)^i (g_1,\ldots,g_i^{(0)},g_i^{(1)},\ldots,g_p \mid g \mid a_0,\ldots,a_q)\\
&+ (-1)^{(p+1)}(g_1,\ldots,g_p, S^{-1}(g^{(2)})(a_0^{(\bar{1})}\ldots a_q^{(\bar{1})})g^{(0)} \mid g^{(1)} \mid a_0^{(\bar{0})},\ldots,a_q^{(\bar{0})})\\
&= \boldsymbol{\delta}(g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q).
\end{aligned}$$

So we have,

$$\begin{aligned}
&\bar{\mathfrak{d}}_0(g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q) = (1,g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q),\\
&\bar{\mathfrak{d}}_i(g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q) = (g_1,\ldots,g_i^{(0)},g_i^{(1)},\ldots,g_p \mid g \mid a_0,\ldots,a_q) \;\; 1\leq i\leq p,\\
&\bar{\mathfrak{d}}_{p+1}(g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q) = (g_1,\ldots,g_p \mid \Delta(g \mid a_0,\ldots,a_q)).
\end{aligned}$$

- To compute $\bar{\mathfrak{t}} = \boldsymbol{\beta}\bar{\tau}\boldsymbol{\gamma}$, we see that,

$$\begin{aligned}
&\bar{\mathfrak{t}}(g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q) = \boldsymbol{\beta}\bar{\tau}(g^{(0)},g^{(1)}\cdot(g_1,\ldots,g_p) \mid a_0,\ldots,a_q)\\
&= \boldsymbol{\beta}(g^{(1)}\cdot(g_1,\ldots,g_p),(a_0^{(\bar{1})}\ldots a_q^{(\bar{1})})g^{(0)} \mid a_0^{(\bar{0})},\ldots,a_q^{(\bar{0})})\\
&= \boldsymbol{\beta}(g^{(1)}g_1,\ldots,g^{(p)}g_p,(a_0^{(\bar{1})}\ldots a_q^{(\bar{1})})g^{(0)} \mid a_0^{(\bar{0})},\ldots,a_q^{(\bar{0})})\\
&= (S^{-1}(g^{(2)}g_1^{(1)})\cdot(g^{(3)}g_2,\ldots,g^{(p+1)}g_p,(a_0^{(\bar{1})}\ldots a_q^{(\bar{1})})g^{(0)}) \mid g^{(1)}g_1^{(0)} \mid a_0^{(\bar{0})},\ldots,a_q^{(\bar{0})})\\
&= (S^{-1}(g_1^{(1)})\cdot(g_2,\ldots,g_p,S^{-1}(g^{(2)})(a_0^{(\bar{1})}\ldots a_q^{(\bar{1})})g^{(0)}) \mid g^{(1)}g_1^{(0)} \mid a_0^{(\bar{0})},\ldots,a_q^{(\bar{0})}).
\end{aligned}$$

So we conclude that

$$\begin{aligned}
&\bar{\mathfrak{t}}(g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q)\\
&= (S^{-1}(g_1^{(1)})\cdot(g_2,\ldots,g_p,S^{-1}(g^{(2)})(a_0^{(\bar{1})}\ldots a_q^{(\bar{1})})g^{(0)}) \mid g^{(1)}g_1^{(0)} \mid a_0^{(\bar{0})},\ldots,a_q^{(\bar{0})}).
\end{aligned}$$

For $\bar{\mathfrak{s}}_i = \boldsymbol{\beta}\bar{\sigma}_i\boldsymbol{\gamma}$, $0\leq i<p$, we have

$$\begin{aligned}
&\bar{\mathfrak{s}}_i(g_1,\ldots,g_p \mid g \mid a_0,\ldots,a_q) = \boldsymbol{\beta}\bar{\sigma}_i(g^{(0)},g^{(1)}\cdot(g_1,\ldots,g_p) \mid a_0,\ldots,a_q)\\
&= \boldsymbol{\beta}(g^{(0)},g^{(1)}\cdot(g_1,\ldots,g_i,g_{i+2},\ldots,g_p) \mid a_0,\ldots,a_q)\epsilon(g_{i+1})\\
&= (g_1,\ldots,g_i,g_{i+2},\ldots,g_p \mid g \mid a_0,\ldots,a_q)\epsilon(g_{i+1}).
\end{aligned}$$

- Next we compute the operator $\mathfrak{b} = \boldsymbol{\beta} b \boldsymbol{\gamma}$. We have

$$\begin{aligned}
&\mathfrak{b}(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = \boldsymbol{\beta} b(g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p) \mid a_0, \ldots, a_q) \\
= \;& \boldsymbol{\beta}(\sum_{0 \le i \le q} (-1)^i (g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p) \mid a_0, \ldots, a_i^{(0)}, a_i^{(1)}, \ldots, a_q) \\
+ \;& (-1)^{q+1} (S^{-1}(a_0^{(0)(\bar{1})}) \cdot (g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p) \mid a_0^{(1)}, a_0, \ldots, a_q, a_0^{(0)(\bar{0})}) \\
= \;& (\sum_{0 \le i \le q} (-1)^i ((S^{-1}(g^{(1)}) g^{(2)}) \cdot (g_1, \ldots, g_p) \mid g^{(0)} \mid a_0, \ldots, a_i^{(0)}, a_i^{(1)}, \ldots, a_q) \\
+ \;& (-1)^{q+1} ((S^{-1}(g^{(1)}) a_0^{(0)(\bar{2})} S^{-1}(a_0^{(0)(\bar{3})})) \cdot (g^{(2)} \cdot (g_1, \ldots, g_p)) \mid S^{-1}(a_0^{(0)(\bar{1})}) g^{(0)} \\
&\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \mid a_0^{(1)}, a_0, \ldots, a_q, a_0^{(0)(\bar{0})})
\end{aligned}$$

$$\begin{aligned}
= \;& (\sum_{0 \le i \le q} (-1)^i (g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_i^{(0)}, a_i^{(1)}, \ldots, a_q) \\
+ \;& (-1)^{q+1} (g_1, \ldots, g_p \mid S^{-1}(a_0^{(0)(\bar{1})}) g \mid a_0^{(1)}, a_0, \ldots, a_q, a_0^{(0)(\bar{0})}).
\end{aligned}$$

So we conclude that,

$$\begin{aligned}
&\mathfrak{d}_i(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = (g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_i^{(0)}, a_i^{(1)}, \ldots, a_q), \;\; 0 \le i \le q, \\
&\mathfrak{d}_{q+1}(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = (g_1, \ldots, g_p \mid S^{-1}(a_0^{(0)(\bar{1})}) g \mid a_0^{(1)}, a_0, \ldots, a_q, a_0^{(0)(\bar{0})}).
\end{aligned}$$

- We consider $\mathfrak{s}_i = \boldsymbol{\beta} \sigma_i \boldsymbol{\gamma}$ and $\mathfrak{t} = \boldsymbol{\beta} \tau \boldsymbol{\gamma}$.

$$\begin{aligned}
&\mathfrak{s}_i(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) \\
&\qquad = \;\; \boldsymbol{\beta}(g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p) \mid a_0, \ldots, a_i, a_{i+2}, \ldots, a_q) \epsilon(a_{i+1}) \\
&\qquad = \;\; (g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_i, a_{i+2}, \ldots, a_q) \epsilon(a_{i+1}).
\end{aligned}$$

Also, for $\mathfrak{t} = \boldsymbol{\beta}\tau\boldsymbol{\gamma}$, we have

$$\begin{aligned}
& \mathfrak{t}(g_1, \ldots, g_p \mid g \mid a_0 . \ldots, a_q) \\
= \; & \boldsymbol{\beta}\tau(g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p) \mid a_0, \ldots, a_q) \\
= \; & \boldsymbol{\beta}(S^{-1}(a_0^{(\bar{1})}) \cdot (g^{(0)}, g^{(1)} \cdot (g_1, \ldots, g_p)) \mid a_1, \ldots, a_q, a_0^{(\bar{0})})) \\
= \; & \boldsymbol{\beta}(S^{-1}(a_0^{(\bar{1})}) g^{(0)}, S^{-1}(a_0^{(\bar{2})}) \cdot (g^{(1)} \cdot (g_1, \ldots, g_p)) \mid a_1, \ldots, a_q, a_0^{(\bar{0})})) \\
= \; & ((S^{-1}(g^{(1)}) a_0^{(\bar{2})} S^{-1}(a_0^{(\bar{3})})) \cdot (g_1, \ldots, g_p)) \mid S^{-1}(a_0^{(\bar{1})}) g^{(0)} \mid a_1, \ldots, a_q, a_0^{(\bar{0})})) \\
= \; & ((S^{-1}(g^{(1)}) \cdot (g_1, \ldots, g_p)) \mid S^{-1}(a_0^{(\bar{1})}) g^{(0)} \mid a_1, \ldots, a_q, a_0^{(\bar{0})})).
\end{aligned}$$

By the above computations, we can state the following theorems.

Theorem 5.1. *The complex* $(\mathsf{C}(C\natural\mathcal{H})_{p,q} \boxtimes \mathsf{W}, b + \mathbf{u}B, \bar{b} + \mathbf{u}\bar{B})$ *is isomorphic to the complex* $(\mathsf{C}^p(\mathcal{H}, \mathsf{C}^q(C^\natural_{\mathcal{H}} \boxtimes \mathsf{W}), \mathfrak{b} + \mathbf{u}\mathfrak{B}, \bar{\mathfrak{b}} + \mathbf{u}\bar{\mathfrak{B}})$, *where* $\bar{\mathfrak{b}}$ *is the Hopf-comodule coboundary on* $\mathsf{C}^q(C^\natural_{\mathcal{H}})$.

Theorem 5.2. *For* $p \geq 0$, $H^p(\mathcal{H}, \mathsf{C}^q(C^\natural_{\mathcal{H}}))$, *with the operators defined as follows, are cocyclic modules, where* $H^0(\mathcal{H}, \mathsf{C}^q(C^\natural_{\mathcal{H}})) \cong \mathsf{C}^\bullet_{\mathcal{H}}(C)$,

$$\begin{aligned}
& \mathfrak{t}(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = ((S^{-1}(g^{(1)}) \cdot (g_1, \ldots, g_p)) \mid S^{-1}(a_0^{(\bar{1})}) g^{(0)} \mid a_1, \ldots, a_q, a_0^{(\bar{0})})), \\
& \mathfrak{d}_i(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = (g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_i^{(0)}, a_i^{(1)}, \ldots, a_q), 0 \leq i \leq q, \\
& \mathfrak{d}_{q+1}(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = (g_1, \ldots, g_p \mid S^{-1}(a_0^{(0)(\bar{1})}) g \mid a_0^{(1)}, a_0, \ldots, a_q, a_0^{(0)(\bar{0})}), \\
& \mathfrak{s}_i(g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_q) = (g_1, \ldots, g_p \mid g \mid a_0, \ldots, a_i, a_{i+2}, \ldots, a_q)\epsilon(a_{i+1}).
\end{aligned}$$

Proof. We prove this for $p = 0$, the general case being similar.

The operators are well defined on the coinvariant subspaces. For example, for $\mathfrak{t}$, since $\mathfrak{t}\bar{\mathfrak{b}} = \bar{\mathfrak{b}}\mathfrak{t}$ and $\bar{\mathfrak{b}}(g \mid a_0, \ldots, a_n) = \boldsymbol{\Delta}(g \mid a_0, \ldots, a_n) - (1 \mid g \mid a_0, \ldots, a_n)$, if $\mathfrak{t}\bar{\mathfrak{b}} = 0$ then $\bar{\mathfrak{b}}\mathfrak{t} = 0$.
To show for example $\mathfrak{t}^{n+1} = 1$, we see that in the coinvariant subspace we have,

$$(1 \mid g \mid a_0, \ldots, a_n) \equiv (S^{-1}(g^{(2)})(a_0^{(\bar{1})} \ldots a_q^{(\bar{1})}) g^{(0)} \mid g^{(1)} \mid a_0^{(\bar{0})}, \ldots, a_q^{(\bar{0})}).$$

So we have,

$$\begin{aligned}
&\bar{\mathfrak{s}}_0 \circ \bar{\mathfrak{t}}(1 \mid g \mid a_0, \dots, a_n) = \bar{\mathfrak{s}}_0 \circ \bar{\mathfrak{t}}(S^{-1}(g^{(2)})(a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g^{(0)} \mid g^{(1)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\\
= \;& \bar{\mathfrak{s}}_0(S^{-1}(g^{(2)})(a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g^{(0)} \mid g^{(1)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\\
= \;& (g^{(1)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\epsilon(S^{-1}(g^{(2)})(a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g^{(0)})\\
= \;& (g \mid a_0, \dots, a_n).
\end{aligned}$$

Therefore,

$$\begin{aligned}
&\bar{\mathfrak{s}}_0 \circ \bar{\mathfrak{t}}(S^{-1}(g^{(2)})(a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g^{(0)} \mid g^{(1)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})})\\
= \;& \bar{\mathfrak{s}}_0(S^{-1}(g^{(3)})(a_0^{(\bar{1})} \dots a_n^{(\bar{1})})g^{(1)} \mid g^{(2)}S^{-1}(g^{(4)})(a_0^{(\bar{2})} \dots a_n^{(\bar{2})})g^{(0)} \mid a_0^{(\bar{0})}, \dots, a_0^{(\bar{0})})\\
= \;& (g^{(2)}S^{-1}(g^{(4)})(a_0^{(\bar{2})} \dots a_n^{(\bar{2})})g^{(0)} \mid a_0^{(\bar{0})}, \dots, a_0^{(\bar{0})})\epsilon(S^{-1}(g^{(3)})(a_0^{(\bar{1})} \dots a_n^{(\bar{1})})g^{(1)})\\
= \;& ((a_0^{(\bar{1})} \dots a_n^{(\bar{1})})g \mid a_0^{(\bar{0})}, \dots, a_0^{(\bar{0})}).
\end{aligned}$$

Therefore, in the coinvariant subspace we have

$$(g \mid a_0, \dots, a_n) = ((a_0^{(\bar{1})} \dots a_n^{(\bar{1})})g \mid a_0^{(\bar{0})}, \dots, a_n^{(\bar{0})}),$$

and then we conclude that

$$\begin{aligned}
&\mathfrak{t}^{n+1}(g \mid a_0, \dots, a_n)\\
= \;& \mathfrak{t}^{n+1}((a_0^{(\bar{1})} \dots a_n^{(\bar{1})})g \mid a_0^{(\bar{0})}, \dots, a_n^{(\bar{0})})\\
= \;& (S^{-1}(a_0^{(\bar{1})} \dots a_n^{(\bar{1})})(a_0^{(\bar{2})} \dots a_n^{(\bar{2})})g \mid a_0^{(\bar{0})}, \dots, a_n^{(\bar{0})})\\
= \;& (g \mid a_0, \dots, a_n).
\end{aligned}$$

Also when $p = 0$, then

$$\begin{aligned}
&\boldsymbol{\delta_0}(g \mid a_0, \dots, a_n)\\
= \;& (1 \mid g \mid a_0, \dots, a_n) - \boldsymbol{\Delta}(g \mid a_0, \dots, a_n)\\
= \;& (1 \mid g \mid a_0, \dots, a_n) - (S^{-1}(g^{(2)})(a_0^{(\bar{1})} \dots a_q^{(\bar{1})})g^{(0)} \mid g^{(1)} \mid a_0^{(\bar{0})}, \dots, a_q^{(\bar{0})}).
\end{aligned}$$

So $\ker \boldsymbol{\delta_0}$ is the coinvariant subspace. $\square$

To compute the cohomologies of the mixed complex $(Tot^\bullet(\mathsf{C}(C\natural\mathcal{H}), b + \bar{b} + \mathbf{u}(B + \bar{B}))$, we filter the complex $\mathsf{C}^\bullet(C\natural\mathcal{H})$ by the subspaces

$$\mathsf{F}^i_{pq} Tot^\bullet(\mathsf{C}(C\natural\mathcal{H}) \boxtimes \mathsf{W}) = \sum_{q \leq i} (\mathcal{H}^{\otimes(p+1)} \otimes C^{\otimes(q+1)}) \boxtimes \mathsf{W}.$$

If we separate the operator $b+\bar{b}+\mathbf{u}(B+T\bar{B})$ as $\bar{b}+(b+\mathbf{u}B)+\mathbf{u}T\bar{B}$. From Theorem(5.1) and (5.2) we can deduce the following theorem.

Theorem 5.3. *The* E^0*-term of the spectral sequence is isomorphic to the complex*

$$\mathsf{E}^0_{pq}=(\mathsf{C}^p(\mathcal{H},\mathsf{C}^q(C^{\natural}_{\mathcal{H}})\boxtimes\mathsf{W}),\boldsymbol{\delta}),$$

and the E^1*-term is*

$$\mathsf{E}^1_{pq}=(H^p(\mathcal{H},\mathsf{C}^q(C^{\natural}_{\mathcal{H}})\boxtimes\mathsf{W}),\mathfrak{b}+\mathbf{u}\mathfrak{B})).$$

The E^2*-term of the spectral sequence is*

$$\mathsf{E}^2_{pq}=HC^q(H^p(\mathcal{H},\mathsf{C}^q(C^{\natural}_{\mathcal{H}}));\mathsf{W}),$$

the cyclic cohomologies of the cocyclic modules $H^p(\mathcal{H},\mathsf{C}^q(C^{\natural}_{\mathcal{H}}))$ *with coefficieints in* W.

We give an application of the above spectral sequence. Let k be a field. Recall that a Hopf algebra $\mathcal{H}$ over k is called cosemisimple if $\mathcal{H}$ is cosemisimple as a coalgebra, that is, every left $\mathcal{H}$-comodule is completely reducible [8]. One knows that a Hopf algebra is cosemisimple if and only if there exists a left integral $x\in\mathcal{H}^{\star}$ with $x(1)=1$ ([8],Theorem 14.0.3). It is easy to see that if $\mathcal{H}$ is cosemisimple and M is an $\mathcal{H}$-bicomodule, then the coalgebra (Hochschild) cohomology groups satisfy $H^i(\mathcal{H},M)=0$ for $i>0$, and $H^0(\mathcal{H},M)=M^{co\mathcal{H}}$, the subspace of coinvariants of the bicomodule M. In fact, we have the following homotopy operator $h:\mathcal{H}^{\otimes n}\otimes M\to\mathcal{H}^{\otimes(n-1)}\otimes M, n\geq 1$,

$$h(g_1,\dots,g_n,m)=x(g_1)(g_2,\dots,g_n,m).$$

One can check theat $\delta h+h\delta=id$. Note that the antipode of $\mathcal{H}$ is bijective if $\mathcal{H}$ is cosemisimple.

Proposition 5.1. *Let* $\mathcal{H}$ *be a cosemisimple Hopf algebra. Then there is a natural isomorphism of cyclic and Hochschild cohomology groups*

$$HC^{\bullet}(C\blacktriangleright\!\!\!<\mathcal{H})\simeq HC^{\bullet}(\mathsf{C}^{\bullet}_{\mathcal{H}}(C)),$$

$$HH^{\bullet}(C\blacktriangleright\!\!\!<\mathcal{H})\simeq HH^{\bullet}(\mathsf{C}^{\bullet}_{\mathcal{H}}(C)).$$

Proof. Since $\mathcal{H}$ is cosemisimple, we have $\mathsf{E}^2_{p,q}=0$ for $p>0$ and the spectral sequence collapses. The first column of E^2 is exactly $H^{\bullet}(\mathcal{H},C^{\natural}_{\mathcal{H}})=\mathsf{C}^{\bullet}_{\mathcal{H}}(C)$. ☐

References

[1] R. Akbarpour, M. Khalkhali, *Hopf algebra equivariant cyclic homology and cyclic homology of crossed product algebras,* math.KT/0011248 v2, (2001).

[2] S. Caenepeel, S. Dăscălescu and S. Rainau, *Cosemisimple Hopf algebras coacting on coalgebras,* Comm. Algebra 24, 1649-1678 (1996).

[3] A. Connes, *Noncommutative Geometry* , Academic Press (1994).

[4] B.L. Feigin and B. L. Tsygan, *Additive K-theory,* K-Theory, Arithmetic and Geometry, Lect. Notes Math. 1289, 67-209, (1986).

[5] E. Getzler and J. D. S. Jones, *The cyclic homology of crossed product algebras*, J. reine angew. Math. 445, 163-174, (1993).

[6] J. L. Loday, *Cyclic Homology,* Springer-Verlag, (1992).

[7] S. Majid, *Foundations of Quantum Group Theory,* Cambridge University Press, (1995).

[8] M. E. Sweedler, *Hopf Algebras,* Benjamin, New York, (1969).

[9] C. Weibel, *An introduction to homological algebra,* Cambridge University Press, (1994).